

## Approach to the Girl with Early Onset of Pubic Hair

Sharon E. Oberfield, Aviva B. Sopher, and Adrienne T. Gerken

Division of Pediatric Endocrinology, Diabetes, and Metabolism, Department of Pediatrics (S.E.O., A.B.S., A.T.G.), Columbia University Medical Center, New York, New York 10032

### Accreditation and Credit Designation Statements

The Endocrine Society is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians. The Endocrine Society has achieved Accreditation with Commendation.

The Endocrine Society designates this Journal-based CME activity for a maximum of 1 AMA PRA Category 1 Credit™. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

### Learning Objectives

Upon completion of this educational activity, participants should be able to

- Evaluate a young girl who presents with early onset of pubic hair
- Manage the clinical care of a child with premature adrenarche
- Describe the progression of metabolic abnormalities in some girls with premature adrenarche

### Target Audience

This Journal-based CME activity should be of substantial interest to endocrinologists.

### Disclosure Policy

Authors, editors, and Endocrine Society staff involved in planning this CME activity are required to disclose to learners any relevant financial relationship(s) that have occurred within the last 12 months with any commercial interest(s) whose products or services are discussed in the CME content. The Endocrine Society has reviewed all disclosures and resolved or managed all identified conflicts of interest, as applicable.

Disclosures for JCEM Editors are found at [http://www.endo-society.org/journals/Other/faculty\\_jcem.cfm](http://www.endo-society.org/journals/Other/faculty_jcem.cfm).

The following individuals reported NO relevant financial relationships:

Sharon E. Oberfield, M.D., Aviva B. Sopher, M.D., Adrienne T. Gerken, A.B., and Leonard Wartofsky, M.D., reported no relevant financial relationships.

Endocrine Society staff associated with the development of content for this activity reported no relevant financial relationships.

### Acknowledgement of Commercial Support

This activity is not supported by grants, other funds, or in-kind contributions from commercial supporters.

### Privacy and Confidentiality Statement

The Endocrine Society will record learner's personal information as provided on CME evaluations to allow for issuance and tracking of CME certificates. No individual performance data or any other personal information collected from evaluations will be shared with third parties.

### Method of Participation

This Journal-based CME activity is available in print and online as full text HTML and as a PDF that can be viewed and/or printed using Adobe Acrobat Reader. To receive CME credit, participants should review the learning objectives and disclosure information; read the article and reflect on its content; then go to <http://jcem.endojournals.org> and find the article, click on CME for Readers, and follow the instructions to access and complete the post-activity test questions and evaluation. The estimated time to complete this activity, including review of material, is 1 hour. If you have questions about this CME activity, please direct them to [education@endo-society.org](mailto:education@endo-society.org).

Activity release date: June 2011

Activity expiration date: June 2012

Premature pubarche, or the development of pubic hair before the age of 8 in girls or 9 in boys, is most commonly caused by premature adrenarche. Adrenarche is the maturation of the adrenal zona reticularis in both boys and girls, resulting in the development of pubic hair, axillary hair, and adult apocrine body odor. Although originally thought to be a benign variant of normal development, premature adrenarche has been associated with insulin resistance and the later development of metabolic syndrome and polycystic ovary syndrome. Although further studies are needed to confirm these relationships, the case presented herein argues for periodic assessment of children at risk. Indeed, recognition of these associations may allow for early preventive measures. (*J Clin Endocrinol Metab* 96: 1610–1622, 2011)

**A**drenarche is the normal maturation of the zona reticularis (ZR) in both boys and girls, resulting in the development of pubic hair, axillary hair, and adult apocrine body odor. These physical changes are preceded by biochemical adrenarche, which has been described to begin physiologically as early as ages 5–6 yr (1) and consists of increased ZR production of  $\Delta 5$  steroids, principally dehydroepiandrosterone (DHEA) and DHEA sulfate (DHEAS).

Premature adrenarche (PA) occurs when signs of adrenarche begin before the age of 8 yr in girls or 9 in boys. The majority of children with PA have idiopathic premature adrenal androgen secretion. Idiopathic PA occurs more frequently in girls than boys by a ratio of about 9:1 (2–5).

The clinical presentation of idiopathic PA is similar to normally timed adrenarche, and adrenal androgens—more correctly termed androgen precursors (6)—principally DHEA, as well as DHEAS, are elevated for chronologic age

Abbreviations:  $\Delta 4$ -A,  $\Delta 4$ -Androstenedione; BMI, body mass index; CAH, congenital adrenal hyperplasia; CPP, central precocious puberty; DEXA, dual-energy x-ray absorptiometry; DHEA, dehydroepiandrosterone; DHEAS, DHEA sulfate; FGIR, fasting glucose-to-insulin ratio; hCG, human chorionic gonadotropin;  $3\beta$ HSD2,  $3\beta$ -hydroxysteroid dehydrogenase type 2; IGFBP-1, IGF binding protein-1; IHL, intrahepatic lipid; IMCL, intramyocellular lipid content; IR, insulin resistance; LDL, low-density lipoprotein; MeS, metabolic syndrome; NC-CAH, nonclassical CAH; oGTT, oral glucose tolerance test; 17OHP, 17-hydroxyprogesterone; PA, premature adrenarche; PCOS, polycystic ovary syndrome; POR, P450 oxidoreductase; PP, premature pubarche; VAT, visceral adipose tissue; ZR, zona reticularis.

but are usually consistent with early puberty or Tanner II–III pubic hair. Smaller elevations are noted for testosterone and  $\Delta 4$ -androstenedione ( $\Delta 4$ -A). There may be modestly increased growth velocity and skeletal age (consistent with height age), but pubertal timing and adult height are usually unaltered (4, 5, 7–9).

Although PA was formerly thought to be a variant of normally timed adrenarche, studies support an association between PA and a history of being small for gestational age and having low birth weight (10), as well as an association with obesity (11). In girls with PA and a history of low birth weight, a 3-fold increase in menarche before age 12 yr has been reported (12).

PA is also associated with a risk of polycystic ovary syndrome (PCOS) and metabolic syndrome (MeS). Metabolic abnormalities reported in prepubertal children with PA include insulin resistance (IR) (13–15), hyperinsulinism (16), increased free IGF-1 (15, 17) and plasminogen activator inhibitor 1 (18), and lower IGF binding protein-1 (IGFBP-1) (15, 16). It has been suggested that in prepubertal girls with PA, plasminogen activator inhibitor 1 levels can predict progression to PCOS (18).

Despite these preliminary data, it is not known which children with PA are most at risk of developing MeS or PCOS. Thus, although PA is frequently a benign process, the diagnosis must carry a heightened suspicion for future metabolic and endocrine abnormalities.

The following patient description will serve to outline the evaluation of the young girl with early onset of pubic hair. We will describe clinical, biochemical, anthropometric, and body composition measures over the period of time from early childhood to late adolescence in a girl with PA. We will also describe management issues, including current controversies regarding treatment modalities.

## Case History

### Age 5 $\frac{4}{12}$ yr

EA (“Early Adrenarche”) is a 5  $\frac{4}{12}$ -yr-old Caucasian girl referred for evaluation of apocrine body odor. She was born full-term and weighed 7 pounds [3175 g; ~30th percentile for sex (19)] at birth. There were no pregnancy complications, including gestational diabetes, and the 17-hydroxyprogesterone (17OHP) newborn screen for congenital adrenal hyperplasia (CAH) was normal. She has grown consistently along the 75th percentile for height, corresponding to the 40th percentile for midparental sex-adjusted target height, but has recently increased her growth velocity to 3.2 cm in 4 months. She has previously plotted in the 50th–75th percentile for weight, but in the past 4 months has plotted in the 75th–90th percentile (Fig.

1). She has not been exposed to lice treatments (tea tree oils), lavender oil, or estrogen sprays or creams and has not increased her intake of soy.

EA’s parents are both of Ashkenazi Jewish heritage. Her mother is 5 feet, 7 inches tall (170.2 cm) and her father is 6 feet tall (183 cm). Midparental sex-adjusted target height is 5 feet, 7 inches. Her mother had menarche at age 12  $\frac{6}{12}$  yr. The patient’s father was treated for severe acne at age 14 yr, had his growth spurt at 15 yr, and developed frontal balding at 30 yr. EA’s 12-yr-old sister has just reached menarche, and her 9-yr-old brother is prepubertal. The patient’s review of systems is negative for excessive thirst or urination, history of diabetes, or headaches.

On examination, EA’s height is 116 cm ( $\approx$ 90th percentile), weight is 22.5 kg ( $\approx$ 90th percentile), and body mass index (BMI) is 16.7 kg/m<sup>2</sup> (83rd percentile). Her blood pressure is 106/70 mm Hg. Funduscopic and thyroid exams are normal. She has no glandular breast tissue (Tanner I). She has faint pubic down on the mons pubis, with sparse curly hairs along the labia majora (Tanner II) and perirectal hairs. Her clitoris is 0.5 cm by 0.3 cm, her posterior labia minora are not fused, and her vagina mucosa is shiny, pink-red, and nonestrogenized. She has no axillary hair, but does have adult-like apocrine odor. Her neurological exam is grossly normal, and she has no areas of skin hyper- or hypopigmentation.

The results of the patient’s initial evaluation (Table 1) include a bone age of 5  $\frac{10}{12}$  to 6  $\frac{10}{12}$  yr, consistent with her height age. Blood work drawn at 0730 h reveals 17OHP of 90 ng/dl,  $\Delta 4$ -A of 66 ng/dl, DHEA of 120 ng/dl, DHEAS of 80  $\mu$ g/dl, testosterone of 8 ng/dl, FSH of 1 mIU/ml, and LH of 0.12 mIU/ml. Fasting serum lipids demonstrate elevated triglycerides, a fasting glucose-to-insulin ratio (FGIR) of 6.5 [normal, < 7 (20)], fasting glucose of 98 mg/dl, and fasting insulin of 15  $\mu$ IU/ml.

### Age 10 $\frac{9}{12}$ yr

EA has been seen every 6 months by her pediatrician and has continued to grow along the 75th–90th percentile for height and 90th percentile for weight. Her pubic hair was Tanner stage III–IV by age 9 yr. Tanner stage II breast tissue was noted at age 9  $\frac{6}{12}$  yr. She had menarche this week and has again been referred to you.

At this visit, EA’s height is 150 cm ( $\approx$ 90th percentile), weight is 48 kg ( $\approx$ 90th percentile), BMI is 21.3 kg/m<sup>2</sup> ( $\approx$ 90th percentile), waist circumference is 70 cm [ $\approx$ 75th percentile (21)], and blood pressure is 110/70 mm Hg. She has a fine moustache, greasy facial skin with comedones, and two pustules on her chin. She has Tanner IV breasts and pubic hair, with mild periareolar, periumbilical, and perirectal hair. The examination is otherwise normal.




FIG. 1. Standardized growth curve for E.A.

EA's height age is 11 1/2 yr, her bone age is 12 1/2 yr, and her estimated adult height is 64.5 ± 2 inches according to the tables of Bailey and Pinneau, which is the 15th percentile for midparental sex-adjusted target height. Her fasting lipid profile demonstrates dyslipidemia (Table 1). Her FGIR has decreased to 4.5, and her oral glucose tolerance test (oGTT) reflects moderate hyperinsulinemia

(22). She has high-normal adrenal and ovarian androgens for Tanner stage with normal gonadotropins.

The patient chooses to participate in an Institutional Review Board-approved body composition protocol. Dual-energy x-ray absorptiometry (DEXA) reveals 35% body fat [≈98th percentile for age, sex, and ethnicity (23)] with increased trunk fat. Measures obtained by magnetic

**TABLE 1.** Clinical and laboratory data by age for EA

|   | Age (yr)                   |                            |                            |
|---|----------------------------|----------------------------|----------------------------|
|   | 5 ¼                        | 10 ½                       | 16 ½                       |
| Height (cm)                                 | 116                        | 150                        | 165.2                      |
| Weight (kg)                                 | 22.5                       | 48.0                       | 77.5                       |
| BMI, kg/m <sup>2</sup> (percentile for age) | 16.7 (83rd %)              | 21.3 (88th %)              | 28.4 (93rd %)              |
| Bone age (yr)                               | 5 ½ to 6 ½                 | 12 ½                       |                            |
| Height age (yr)                             | 6 ¼                        | 11 ¾                       |                            |
| Predicted height (cm)                       |                            | 164                        |                            |
| Waist circumference (cm)                    | 54 (50–75th %)             | 70 (75th %)                | 84 (75th %)                |
| Breasts                                     | Tanner I                   | Tanner IV                  | Tanner IV                  |
| Pubic hair                                  | Tanner II                  | Tanner IV                  | Tanner IV                  |
| Ferriman-Gallwey score                      |                            |                            | 16                         |
| Blood pressure (mm Hg)                      | 106/70                     | 110/70                     | 118/76                     |
| Testosterone (ng/dl)                        | 8 (7–28)                   | 28 (13–32)                 | 41 (13–32)                 |
| Δ4 (ng/dl)                                  | 66 (<10–72)                | 194 (47–208)               | 306 (47–208)               |
| 17OHP (ng/dl)                               | 90 (11–98)                 | 114 (18–230)               | 121 (18–230)               |
| DHEAS (μg/dl)                               | 80 (34–129)                | 237 (58–260)               | 282 (58–260)               |
| LH (mIU/ml)                                 | 0.12 (0.02–4.7)            | 4.4 (0.4–11.7)             | 19.0 (0.4–11.7)            |
| FSH (mIU/ml)                                | 1.0 (1.0–10.8)             | 2.1 (1.5–11.7)             | 5.4 (1.5–11.7)             |
| Total cholesterol (mg/dl)                   | 140 (126–205) <sup>a</sup> | 165 (124–201) <sup>a</sup> | 174 (120–200) <sup>a</sup> |
| LDL (mg/dl)                                 | 89 (68–140) <sup>a</sup>   | 112 (68–136) <sup>a</sup>  | 126 (60–135) <sup>a</sup>  |
| HDL (mg/dl)                                 | 45 (36–73) <sup>a</sup>    | 36 (37–70) <sup>a</sup>    | 34 (35–73) <sup>a</sup>    |
| Triglycerides (mg/dl)                       | 144 (32–105) <sup>a</sup>  | 162 (37–131) <sup>a</sup>  | 190 (39–124) <sup>a</sup>  |
| Fasting glucose (mg/dl)                     | 98                         | 100                        | 94                         |
| Fasting insulin (μIU/ml)                    | 15                         | 22                         | 18                         |
| FGIR  | 6.5 (<7) <sup>b</sup>      | 4.5 (<7) <sup>b</sup>      | 5.2 (<7) <sup>b</sup>      |
| oGTT <sup>c</sup>                           |                            |                            |                            |
| 0 min                                       | 98/15                      | 100/22                     | 94/18                      |
| 30 min                                      | 138/54                     | 165/92                     | 156/70                     |
| 60 min                                      | 102/18                     | 141/86                     | 144/62                     |
| 90 min                                      | 108/40                     | 117/58                     | 138/52                     |
| 120 min                                     | 112/38                     | 118/66                     | 116/36                     |

Normal values for sex and age are shown in *parentheses*. HDL, High-density lipoprotein.

<sup>a</sup> Represents 5th to 95th percentile values for respective age ranges (103).

<sup>b</sup> From Ref. 20.

<sup>c</sup> oGTT data are expressed as glucose (mg/dl)/insulin (μIU/ml).

resonance spectroscopy for intramyocellular lipid content (IMCL) and intrahepatic lipid (IHL) content are increased compared with her age-matched peers.

### Age 16 ½ yr


EA has been seen two times per year for the past 6 yr. After menarche, she had irregular cycles (every 2–4 months). Transabdominal ultrasound of the pelvis showed 4-cc ovaries with multiple peripheral follicles. At the age of 12 ½ yr, after 2 yr of persistently irregular cycles and increasing hyperinsulinemia, she was started on extended release metformin at a dose of 500 mg/d for 1 wk and increased to 1000 mg/d thereafter. Menses now occur every 4–5 wk.

EA has followed a Mediterranean diet (low glycemic index, low carbohydrates, high grain and fiber content). She does aerobic and resistance exercises three times per week. She reports continued acne, for which she uses topical drying agents. She is upset about her body and facial

hair and regularly undergoes laser hair removal with significant improvement.

Her height is 165.2 cm (50th–75th percentile), weight is 77.5 kg (≈95th percentile), BMI is 28.4 kg/m<sup>2</sup> (>90th percentile), waist circumference is 84 cm [75th percentile (21)], and blood pressure is 118/76 mm Hg. She has significant facial hair, periumbilical and perirectal hairs, and mild acne. Her Ferriman-Gallwey score is elevated at 16. She has Tanner IV breasts and pubic hair and faint acanthosis nigricans of the neck and periaxillary areas. Laboratory tests show elevated Δ4-A, DHEAS, and LH and borderline-high fasting lipids. oGTT shows an FGIR of 5.2. She now has 40% body fat on DEXA [>99th percentile for age, sex, and ethnicity (23)].

Dietary and exercise interventions are reviewed and weight loss of 10–15 pounds over the next year is recommended. In preparation for possible initiation of a low-dose oral contraceptive pill containing ethinyl estradiol and norgestimate or norgestrel, which may improve her



**FIG. 2.** Adrenal steroidogenesis.  $17\beta$ HSD1,  $17\beta$  hydroxysteroid dehydrogenase, type 1;  $17\beta$ HSD2,  $17\beta$  hydroxysteroid dehydrogenase, type 2;  $17\beta$ HSD3,  $17\beta$  hydroxysteroid dehydrogenase, type 3;  $3\beta$ HSD,  $3\beta$  hydroxysteroid dehydrogenase; Arom, aromatase; b5, cytochrome b5; DOC, 11-deoxycorticosterone; Isom, isomerase; P450c11 $\beta$ , 11 $\beta$ -hydroxylase; P450c17, 17 $\alpha$ -hydroxylase and 17,20-lyase; P450c18, 18-hydroxylase; P450c21, 21 $\alpha$ -hydroxylase; P450scc, cholesterol side-chain cleavage enzyme (or desmolase); StAR, steroidogenic acute regulatory protein; SULT2A1, dehydroepiandrosterone sulfotransferase.

acne and hirsutism, a negative family history for clotting disorders is confirmed.

## Discussion

EA's course from childhood through midadolescence is representative of a child who presents with PA. Her clinical findings will be discussed in detail after a review of the pathophysiology, evaluation, and clinical spectrum of adrenarche.

### Adrenal development

The early adrenal cortex consists of the fetal zone, which involutes completely by the end of the first year of life (1), and the definitive zone, which persists and compartmentalizes to become the mature adrenal cortex.

The fetal zone is ACTH-dependent (24, 25) and becomes the primary site of fetal steroidogenesis at around 16–20 wk of gestational age. The outer portion of the fetal zone, termed the transitional zone, is the site of fetal cortisol synthesis. The fetal zone also produces DHEA, a precursor for the fetal estrogens required to sustain pregnancy, and DHEAS. The fetal zone involutes rapidly after birth (24, 26), and DHEA and DHEAS levels decline with postnatal regression of the fetal zone and remain low until adrenarche (27).

The definitive zone, or neocortex, begins as a narrow band of cells outside the fetal zone and continues to grow into the postnatal period, when it is remodeled to become

the adult adrenal cortex (25). The adrenal medulla, meanwhile, is populated by neural crest cells concurrent with encapsulation of the adrenal. The neural crest cells differentiate into catecholamine-producing chromaffin cells that persist as medullary islands until birth, when they coalesce to form the immature medulla (27).

Postnatally, the adrenal cortex is remodeled to consist of continuous bands of zona glomerulosa and fasciculata cells. Islands of reticularis cells appear around the age of 3 yr and eventually coalesce to form a continuous ZR at 6–8 yr of age, concurrent with the beginning of biochemical adrenarche. The ZR continues to expand from adrenarche through early adolescence, peaking at around age 13 (1).

### Classical steroid pathways in DHEA and DHEAS synthesis

The biochemical hallmark of adrenarche is increased production of DHEA and DHEAS by the ZR without a concomitant rise in cortisol. DHEA and DHEAS are produced in the ZR via the  $\Delta^5$  pathway of steroidogenesis, and DHEA may be converted to  $\Delta^4$ -A (Fig. 2).

#### CYP11A1 (P450scc)

The first committed step in adrenal steroidogenesis is the conversion of cholesterol to pregnenolone by the cholesterol side-chain cleavage enzyme, CYP11A1 (P450scc), which is expressed throughout the adrenal cortex. CYP11A1 is regulated acutely by the steroidogenic acute regulatory protein, which mediates the transfer of cholesterol into the inner mitochondrial membrane (28, 29) and chronically via transcriptional regulation (29). CYP11A1 expression does not change with adrenarche (30).

#### CYP17 (P450c17)

Qualitative regulation of steroidogenesis occurs via CYP17 (P450c17), which has a dual enzymatic role in the adrenal: 17 $\alpha$ -hydroxylation of pregnenolone and progesterone, and 17,20-lyase activity on their 17-hydroxy derivatives (31). Both the 17 $\alpha$ -hydroxylase and 17,20-lyase actions of CYP17 are required to produce adrenal androgens, and the relative efficiency of these two reactions helps to determine the types of steroids produced. In young children, the 17 $\alpha$ -hydroxylase activity of CYP17 produces cortisol, but 17,20-lyase activity (and thus androgen production) is low. During

adrenarche, an increase in 17,20-lyase activity leads to increased DHEA and DHEAS production (32).

Several factors influence the balance of 17 $\alpha$ -hydroxylase and 17,20-lyase activity of CYP17. The flavoprotein P450 oxidoreductase (POR) transfers electrons from reduced NADPH (nicotinamide adenine dinucleotide phosphate) and is required for both functions of CYP17 (33). *In vitro*, as the ratio of POR to CYP17 increases, 17,20-lyase activity is preferentially enhanced, leading to DHEA production (34). The 17,20-lyase activity of CYP17 is also augmented by cytochrome b5 (b5, CYB5), an allosteric effector of the CYP17-POR interaction (35–37).

Finally, the 17,20-lyase activity of CYP17 is enhanced by phosphorylation of serine and threonine residues by a cAMP-dependent protein kinase (38) and decreased by dephosphorylation via protein phosphatase 2A (39). CYP17 phosphorylation has been suggested as a possible link between the adrenal hyperandrogenism and IR in PA and PCOS (38) because serine phosphorylation of the insulin receptor has been shown *in vitro* to cause IR (40, 41).

### 3 $\beta$ HSD2

The balance of adrenal androgen production is also affected by 3 $\beta$ -hydroxysteroid dehydrogenase type 2 (3 $\beta$ HSD2), which catalyzes the conversion of DHEA to  $\Delta$ 4-A (31, 42). This conversion diverts adrenal androgens from the  $\Delta$ 5 pathway, which produces DHEA and DHEAS, to the  $\Delta$ 4 pathway, which produces  $\Delta$ 4-A. Recent *in vitro* evidence has shown that 3 $\beta$ HSD2 is inhibited by cortisol, raising the possibility that rising intraadrenal cortisol levels may play a role in the initiation of adrenarche (43). Peripheral conversion of  $\Delta$ 4-A produces the potent androgens testosterone and dihydrotestosterone. During adrenarche, expression of 3 $\beta$ HSD2 in the ZR decreases (37, 44), increasing adrenal production of DHEA and DHEAS relative to  $\Delta$ 4-A.

### SULT2A1

The final step in the  $\Delta$ 5 pathway is the conversion of DHEA to DHEAS via DHEA sulfotransferase (SULT2A1) (31, 45). SULT2A1 expression increases during adrenarche in association with maturation of the ZR (37). Sulfation limits the pool of DHEA available for conversion to more potent androgens, and defects in sulfation have been associated with PA (45).

## Clinical findings

### Hyperinsulinemia and IGF-1

Decreased insulin sensitivity is a key component of MeS (46, 47). Numerous studies have demonstrated diminished insulin sensitivity in children with PA (13, 14, 16, 20, 48). Of note, Ibáñez *et al.* (49) have also reported that, in

a population of Spanish girls, first-degree relatives of girls with PA were at greater risk for impaired glucose tolerance and type 2 diabetes.

In a small group of prepubertal girls with PA, Ibáñez *et al.* (48) reported that girls with PA had increased mean serum insulin levels and increased early insulin response to oGTT; however, PA subjects had a higher average BMI than controls. No comparison was made between obese and nonobese PA girls (16, 48). Denburg *et al.* (13) demonstrated decreased insulin sensitivity on oGTT and elevated IGF-1 levels independent of obesity in prepubertal boys with PA, although this was not seen in a cohort of lean Spanish boys (10).

*In vitro*, both IGF-1 and insulin have been shown to potentiate LH-stimulated androgen synthesis in rat (50) and human (51) theca-interstitial cells, suppress SHBG production in hepatoma cell lines (52), and induce steroidogenic enzymes in cultured human adrenocortical cells (25, 53). Additionally, both women with PCOS and patients with PA have decreased levels of IGFBP-1, which are inversely correlated with fasting insulin levels and ACTH-stimulated adrenal steroid levels (16).

Taken together, these findings suggest a link between IR, IGF-1, and the hyperandrogenemia of PA and PCOS (Fig. 3) (17). Some authors have proposed that hyperinsulinemia is the inciting event in a cascade leading to hyperstimulation of ovarian and/or adrenal androgen synthesis, such that in some populations, IR and hyperandrogenism are causally related (53, 54).

### Obesity

Although there appears to be an association between obesity and PA (11), the exact nature of this relationship




FIG. 3. Proposed mechanism for the development of PCOS or MeS in patients with PA. [Adapted from M. E. Silfen *et al.*, *J Clin Endocrinol Metab* 87:398–403, 2002 (17). © The Endocrine Society.]

is unclear. A small study found that obese African-American and Caribbean Hispanic adolescent females with a history of both childhood obesity and PA were hyperandrogenic and hyperinsulinemic (55, 56). The emerging picture based on limited studies of prepubertal Hispanic and African-American girls with PA is that most obese girls with PA have decreased insulin sensitivity compared with nonobese girls with PA (15, 17, 20, 57). Additional data are needed regarding other ethnic groups. Recently, Marshall's group (58) has demonstrated that obesity in the prepubertal and early pubertal child may enhance LH secretion and ovarian androgen production, providing additional evidence linking obesity and hyperandrogenism.

### **PCOS and/or functional ovarian hyperandrogenism**

In a cohort of Catalan girls with a history of PA, nearly half of the postpubertal females with a history of PA had hirsutism, oligomenorrhea, and/or elevated androgen levels (16, 59, 60). Patients with a history of PA and low birth weight also had an increased prevalence of anovulatory cycles (10, 61). However, none of these results were analyzed with respect to the impact of obesity. In contrast, in a cohort of pubertal Finnish females with a history of childhood PA, no hyperandrogenism was observed (9).

It is difficult to assess what proportion of girls with PA progress to PCOS, in part due to multiple consensus statements for the diagnosis of PCOS (62, 63). We have suggested that a definitive diagnosis of PCOS in adolescents should not be made until at least 2 yr after menarche and should require the presence of hyperandrogenism, chronic anovulation, and polycystic ovaries. In addition, we have suggested that hyperandrogenism be defined as hyperandrogenemia and that clinical findings such as acne or alopecia should be discounted, with the exception of documented progressive hirsutism (64).

### **Lipid abnormalities**

Dyslipidemia, including increased triglycerides and low high-density lipoprotein, is a component of MeS in both adults and children and portends future risk of type 2 diabetes (65, 66). Dyslipidemia has been demonstrated in girls with PA and a history of PA, primarily manifested by hypertriglyceridemia (48). Güven *et al.* (67) demonstrated higher mean total cholesterol, low-density lipoprotein (LDL), VLDL (very low-density lipoprotein), and atherogenic index in prepubertal Turkish girls with PA than in controls, and Denburg *et al.* (13) have shown a trend toward higher triglycerides in a BMI-adjusted analysis of prepubertal boys with PA. Although another study showed no difference in lipoprotein (a) levels between prepubertal girls with PA and controls, pubertal girls with a

history of PA had significantly higher lipoprotein (a) levels than their age-matched peers (68).

Although dyslipidemia represents an independent risk factor for cardiovascular disease, an additional question is the extent to which abnormal lipid deposition within the tissues may contribute to IR and type 2 diabetes. Although the relationship between IR and body composition, particularly abdominal fat, IMCL, and IHL, is just beginning to be investigated, preliminary results have yielded interesting hypotheses regarding the origin of IR in these patients. Indeed, the "adipose-tissue expandability hypothesis" argues that IR in obesity and other disorders, including PCOS, is attributable to lipotoxicity due to the inability to expand adipose tissue in a metabolically safe manner (69). The relationship between IR and PA is currently being investigated.

### **Bone mineral density and body composition**

To date, three reports have addressed body composition in exclusively prepubertal populations with PA (70–72). Our group has found increased bone mineral content and bone mineral density using DEXA in prepubertal girls with PA, compared with control subjects (71).

We have also found increased IMCL in PA girls compared with controls, but no difference in visceral adipose tissue (VAT), abdominal SAT (subcutaneous adipose tissue), or VAT:SAT ratio. Of note, however, this population consisted of a very small group of six girls with PA and eight controls (70).

The critical question with respect to body composition in PA is its relationship with IR and risk for MeS. Ibáñez *et al.* (73) have evaluated body composition by DEXA in prepubertal and adolescent girls (aged 6–18 yr) with a history of PA. Although BMI did not differ between PA and control girls, those with PA had greater total and trunk fat mass in the prepubertal and pubertal periods. Furthermore, trunk fat mass showed an independent and significant positive relationship with fasting insulin and free androgen index (73). In a group of adolescent girls and young women with varying degrees of hyperandrogenism, hyperinsulinism, and anovulation, Ibáñez *et al.* (74–77) found that short-term therapy with metformin or a metformin-flutamide combination resulted in decreased fat mass, increased lean body mass, and improved androgens, insulin sensitivity, and serum lipids without a change in overall body mass.

### **Differential diagnosis**

Premature adrenarche must be distinguished from a number of other benign and pathological conditions (Table 2).

**TABLE 2.** Differential diagnosis of early onset of pubic hair

|                            | Pubic hair | Breast development, testicular enlargement, other physical findings                     | Androgens   | LH and FSH       | Bone age                            | Growth velocity | Additional notes  |
|----------------------------|------------|---|---|------------------|-------------------------------------|-----------------|---|
| Isolated PP                | Present    | Absent  | Prepubertal, DHEAS not elevated   | Prepubertal      | ↔                                   | ↔               | Possibly related to increased androgen receptor sensitivity |
| PA                         | Present    | Absent  | ↑ but appropriate for Tanner II-III                                     | Prepubertal      | Slight ↑ correlated with height age | Slight ↑        | Most common cause of PP                                     |
| Precocious puberty         | Present    | + Breast development, + testicular enlargement  | ↑↑ for males, estradiol can be ↑ for females                            | Early pubertal   | ↑                                   | ↑↑              | More common in girls; more often pathological in boys       |
| NC-CAH                     | Present    | +/- Penile/clitoral enlargement, absent breast development or testicular enlargement    | ↑↑ May be elevated for Tanner stage, with ↑ adrenal androgen precursors | Prepubertal      | ↑                                   | ↑               | ↑ 17OHP in 21-hydroxylase deficiency (most common form)     |
| Virilizing tumors          | Present    | +/- Breast development, + penile/clitoral enlargement, infantile testes                 | Usually ↑ but variable, depending on tumor expression                   | Prepubertal or ↓ | ↑                                   | ↑               | May have ↑ DHEA, DHEAS, T, Δ4-A, or βhCG                    |
| Exogenous hormone exposure | Present    | Variable; may have breast development, small testes, and/or penile/clitoral enlargement | Variable, depends on specific exposure                                  | Prepubertal      | ↑                                   | ↑↑              | Take careful medication history from all caretakers         |

T, Testosterone; ↑, increased; ↓, decreased; ↑↑, markedly increased; ↓↓, markedly decreased; ↔, normal for age.

### Premature pubarche without elevated androgens

Although PA is the most common cause of premature (or precocious) pubarche (PP), the latter occasionally occurs in isolation, without a concomitant rise in adrenal androgens. Bone age is not advanced in isolated PP, and growth velocity is not accelerated for age. It has been speculated that these individuals have an increased receptor sensitivity to low circulating levels of androgens (78).

### Precocious puberty

Precocious puberty may be central (also called gonadotropin-dependent precocious puberty) or peripheral (also termed gonadotropin-independent). Central precocious puberty (CPP) is generally complete and may be idiopathic, while peripheral precocious puberty is often incomplete and always pathological in nature. Precocious puberty is more common in girls by a ratio of 5:1, and idiopathic CPP by 9:1 (79). However, in girls 80% of precocious puberty is idiopathic, while approximately two thirds of males with precocious puberty have central nervous system pathology (79, 80).

Idiopathic CPP may present with pubic or axillary hair but is differentiated clinically from PA by the presence of breast or testicular development. It is caused by early maturation of the hypothalamic-pituitary-gonadal axis and is more common in African-American children compared with their Caucasian or Hispanic peers (81, 82). Androgens (and sometimes estradiol) are elevated for age, and gonadotropins are in the early pubertal range. Bone age and growth velocity are accelerated, which may lead to reduced adult height (83, 84). Of note, prior exposure to elevated serum androgens (*e.g.* virilizing tumors, untreated CAH) may result in CPP due to early maturation

of the hypothalamus (85, 86). For a review of precocious puberty, please refer to Carel and Léger's article (87).

### Nonclassical CAH (NC-CAH)

NC-CAH may also present with PP. Estimates of the proportion of cases of PP due to NC-CAH vary widely, from 0 to 40% (53), although these differences may be due to ethnic differences between study populations. NC-CAH is the result of mild deficiencies in the same enzymes responsible for classical CAH, namely 21-hydroxylase, 11-hydroxylase, or 3βHSD2. These deficiencies are typically not sufficient to cause an abnormal newborn screening for CAH; *i.e.* in nonclassic 21-hydroxylase deficiency, basal 17OHP levels shortly after birth are within normal limits. The diagnosis is suspected when androgens are elevated in the presence of prepubertal gonadotropins. Accelerated height velocity and a moderate increase in bone age raise suspicion for NC-CAH, as do marked virilization (clitoral or penile growth) or early development of cystic acne (88). Clinical variables, however, have poor predictive value for distinguishing NC-CAH from idiopathic PA (89).

The diagnosis of NC-CAH is confirmed when elevated levels of the appropriate precursor steroids are noted after ACTH stimulation. However, there is ongoing controversy regarding which children with PP should receive ACTH stimulation testing. Those for whom there is a heightened index of suspicion, and therefore greater indication for testing, include children with elevated basal androgens, bone age advancement greater than 1–2 yr above chronological age, or virilization in the absence of true puberty. Early morning basal 17OHP above 200 ng/dl is 100% sensitive and 99% specific for NC-CAH (89).



### Virilizing tumors

Virilizing adrenal or gonadal tumors are a rare cause of PP. Clinically, these tumors may present with rapidly progressive and/or exaggerated symptoms of virilization, such as clitoral enlargement or rapid penile growth. Growth velocity is notably accelerated, and bone age is advanced and rapidly progressive. Serum androgens are markedly increased, while gonadotropins are prepubertal or suppressed (79). Serum  $\beta$ -human chorionic gonadotropin (hCG), which acts at gonadal LH receptors to stimulate testosterone production, may be elevated in hCG-producing tumors of the testis, brain, or liver.

### Exogenous steroid exposure

PP with feminizing signs may be attributable to estrogen exposure from creams, oils, or sprays (e.g. EvaMist) (90) prescribed for caretakers, or from exposure to lavender or tea tree oil, which have estrogenic activity (91). Synthetic environmental endocrine disruptors such as bisphenol A also appear to have estrogenic effects that may alter the course of normal pubertal development (92), although the mechanisms and extent of these effects are still being explored.

Exogenous androgen exposure from creams or gels containing testosterone or  $\Delta$ 4-A may precipitate PP with other androgenic signs, usually in the absence of elevation of androgen precursors, thus mimicking PA (93). Importantly, it should be noted that exogenous steroids need not be applied directly to the patient but may be transferred via skin-to-skin contact (90, 93).

### Laboratory and imaging assessment

A definitive algorithm for the evaluation of PA cannot be recommended because some variance is due to the clinical concern of the physician. In our practice, all patients with PA receive baseline blood tests for androgens (testosterone,  $\Delta$ 4, DHEAS), gonadotropins (LH and FSH), and 17OHP. Thyroid studies (TSH and free  $T_4$ ) are ordered if hypothyroidism is suspected. If early morning 17OHP is above 200 ng/dl, an ACTH-stimulation test to evaluate for NC-CAH is indicated (89). In boys,  $\beta$ -hCG should be obtained to rule out an hCG-secreting tumor.

The decision to order imaging studies is based on clinical impression and results of initial laboratory evaluation. Rapid virilization or physical signs of precocious puberty are indications to obtain an x-ray of the left hand to determine bone age. If initial studies show very elevated DHEAS, abdominal ultrasound may be indicated as an initial screen to evaluate for adrenal neoplasm. Abdominal computed tomography or magnetic resonance imaging is indicated if a mass is noted, although controversy still exists regarding which of these is preferred. Similarly, ul-

trasonographic evaluation of the ovaries or testes may be indicated if significantly elevated testosterone or  $\Delta$ 4-A is found, if a testicular mass is palpated, or if there is significant testicular asymmetry. Elevated  $\beta$ -hCG in boys should prompt an evaluation for the presence of a tumor of the testis, liver, or brain. Finally, brain magnetic resonance imaging with contrast may be indicated in precocious puberty (especially in boys), or if there is any suspicion of central nervous system pathology.

### Management

Once the diagnosis of PA is made, families should be counseled regarding the increased risk of early puberty and should be given instructions to return promptly if marked growth acceleration or progressive virilization occur. In the majority of cases, when clinical and laboratory findings are consistent with typical PA, these children may be followed by a general pediatrician, family physician, or pediatric endocrinologist at approximately 6-month intervals, with some variation in the frequency of follow-up depending on the preference of the physician. Careful assessment of growth velocity, weight gain, and progression of signs of androgen excess are indicated. Yearly fasting measures of IR (such as FGIR) and lipid levels are recommended, particularly if there has been significant weight gain. Some clinicians follow adrenal androgen levels.

Most children do not demonstrate rapid pubertal progress or bone age advancement and do well with lifestyle interventions. However, should the rate of pubertal changes suddenly accelerate, a bone age and androgen levels (testosterone,  $\Delta$ 4, DHEAS) should be repeated to reassess the child's status. If not previously performed, an ACTH-stimulation test should be conducted to exclude defects of adrenal steroidogenesis.

In our practice, we continue to follow girls with a history of PA at least yearly after menarche. We encourage continued adherence to a low-fat, low-glycemic index diet and regular exercise. If menstrual regularity has not been established 2 yr after menarche, additional evaluation is undertaken to assess ovarian steroids, and a pelvic sonogram is obtained. At that point, we may recommend metformin, particularly if there is evidence of IR, and/or an oral contraceptive pill.

### Areas of current controversy

#### Pharmacotherapy

In pediatrics, medications are usually employed to treat a disease. However, it is not clear whether PA, *per se*, qualifies as a disease, although it often is an antecedent to adolescent and adult metabolic disease, androgen excess, or PCOS. Because PA does not always progress to a state of metabolic dysregulation, the question of how to identify

which young children require pharmacotherapy remains. If we view PA as a potential antecedent of adult disease, earnest thought must be given to the benefit of early intervention with altered lifestyle and/or medical therapy. Because treatment may be continued for years, the long-term risk of pharmacological intervention must also be carefully weighed.

A common factor in the progression of metabolic disturbances associated with PA is IR, a feature shared with PCOS. In PCOS, metformin has been reported to have salutary effects on menstrual irregularity, fertility, weight, and hyperandrogenism (94, 95). However, studies of the effects of metformin treatment on metabolic changes in young patients and populations at risk for PCOS have been limited. Metformin acts through AMP kinase, a regulator of energy metabolism that restores ATP levels by switching on catabolic pathways (glycolysis, fatty acid oxidation) and switching off gluconeogenesis. AMP kinase may also affect reproductive functions including granulosa cell steroidogenesis, nuclear oocyte maturation, and hypothalamic LH secretion and may be involved in adipokine signaling. Other effects of metformin include increased glucose uptake through facilitation of glucose transporter translocation to the plasma membrane, increased insulin receptor tyrosine kinase activity, and reduced availability of IGF-1 through increased IGFBP-1 (95).

Although formal guidelines are not presently available for pharmacotherapy in PA, preliminary studies suggest that in girls with PA and a history of low birth weight and rapidly progressive puberty, metformin may reverse the trends of advancing weight gain, increasing truncal fat mass, and lipid abnormalities (75). Ibáñez *et al.* (96) also used metformin in an effort to halt the advancement of rapidly progressive puberty in girls as young as 8 yr of age. After 4 yr, girls in the treated group were thinner, less insulin resistant, and less hyperandrogenic; had less visceral fat; and had lower IGF-1 levels than untreated girls (96).

A recent 6-yr, randomized, controlled, longitudinal study of metformin treatment in 38 girls with PA and a history of being small for gestational age, who were thus at high risk for development of early menarche and PCOS, demonstrated the lasting beneficial effects of metformin for at least 1 yr after termination of treatment. Adolescents who were previously treated with metformin had later menarche compared with controls, accompanied by a more normal distribution of age at menarche. More than 2 yr after menarche, treated adolescents had less IHL and VAT, greater lean mass, and lower testosterone and fasting insulin levels compared with controls (97). During treatment, IGF-1 levels were significantly lower in the

treatment group (98), but treated adolescents were closer to their target heights at 2 yr after menarche (97).

Despite promising results of metformin therapy, longer-term studies in larger populations are needed before universal recommendations for drug therapy can be made.

## Back to the Patient

### Age 5 $\frac{4}{12}$ yr

At the time of presentation, EA had signs of androgen excess, *i.e.* growth acceleration and onset of body odor and pubic hair. Although she had a normal newborn screen for 21-hydroxylase deficiency, it is not usually positive for patients with NC-CAH. Additionally, EA's family history may suggest NC-CAH, given her Ashkenazi ancestry and her father's history of acne and early frontal balding. However, EA's normal 17OHP on initial evaluation was not suggestive of NC-CAH. Her negative history of exposure to steroid-containing compounds excludes an exogenous source of androgens or estrogens.

On physical examination, the absence of clitoromegaly supported the diagnosis of PA rather than a pathological virilizing process. Lack of breast tissue and prepubertal gonadotropins were not consistent with early puberty. Although her parents were tall, her increase in growth velocity was suspicious. However, her bone age was not significantly advanced, and her adrenal androgens were concordant with her Tanner II stage of pubic hair development. These findings exclude an adrenal tumor as the cause of her PP. It is noteworthy that even at this young age EA demonstrated elevated triglycerides and a slight decrease in her FGIR suggestive of early IR, which could be consistent with the adipose-tissue expandability hypothesis (69).

### Age 10 $\frac{9}{12}$ yr

EA had onset of breast tissue at a normal age but had relatively early menarche. Mean age of menarche for Caucasian females is close to 12 yr (81, 99). At this time, she was overweight and had excess facial and body hair and comedones. Hirsutism may be associated with adrenal hyperandrogenism or may be an early manifestation of PCOS.

With bone age advancement, EA's height prediction was in the lower range for her midparental sex-adjusted target height, although she was still within the normal range for adult female height. Idiopathic PA is usually not associated with a significantly diminished adult height (4, 5).

EA's elevated trunk fat, percentage body fat, IMCL, and IHL are all consistent with a picture of developing PCOS and MeS. Ectopic fat deposition refers to accumulation of fat in the skeletal muscle (IMCL), liver (IHL), and

trunk (VAT). The progression of IR likely involves ectopic fat deposition in the following sequence: IMCL leads to skeletal muscle IR, followed by dyslipidemia, IHL deposition, hepatic IR, VAT accumulation, and intrapancreatic lipid deposition. This causes the abnormal release of adipokines, resulting in the development of MeS and type 2 diabetes (100). Our preliminary data from nonobese adolescents with PCOS and controls show that IMCL correlates with measures of IR, while IHL correlates with dyslipidemia, supporting this proposed sequence (101).

### Age 16 <sup>6</sup>/<sub>12</sub> yr

EA was counseled on diet and exercise to decrease her relative IR. Two years after the onset of menses, she was started on metformin for persistently irregular cycles and hyperinsulinemia. Although limited long-term data are available on outcomes of girls with PA treated with metformin, this appears to be a reasonable intervention in an attempt to decrease IR. Despite metformin use, she continued to have borderline elevation of her lipids and blood glucose.

During the progression of puberty, her weight percentile stabilized and she did not become hypertensive, although her hormone levels were consistent with ovarian hyperandrogenism or PCOS. The issue of use of oral contraceptive pills in this population is somewhat controversial because they can aggravate low-grade inflammation and adiposity (102), but oral contraceptive pills can serve to decrease free circulating androgens by increasing SHBG levels and thus result in improvement of acne.

### Conclusion

EA's history underscores the need for careful clinical and laboratory follow-up of children with PA. Although many children who present with PA will not have substantial metabolic risk, some indeed are at risk for adult MeS and PCOS. The task of the future will be to identify those at greatest risk and discern who might benefit from both lifestyle changes and early pharmacological intervention. If the inciting event in the cascade leading to PA is, in fact, hyperinsulinemia, efforts to reverse this process with agents such as metformin may prove reasonable.

### Acknowledgments

Address all correspondence and requests for reprints to: Sharon E. Oberfield, M.D., 622 West 168th Street, PH 5E-522, New York, New York 10032. E-mail: seo8@columbia.edu.

This work was supported by National Institutes of Health (NIH)/National Institute of Diabetes and Digestive and Kid-

ney Diseases (NIDDK), Training Grant in Pediatric Endocrinology T32 DK065522 (Principal Investigator, S.E.O.); NIH/NIDDK Medical Student Research Training Program Grant 3T32DK065522-06S1 (to A.T.G.); and a Doris Duke Clinical Research Foundation grant to Columbia University College of Physicians and Surgeons (to A.T.G.).

Disclosure Summary: The authors have nothing to disclose.

### References

- Dhom G 1973 The prepubertal and pubertal growth of the adrenal (adrenarche). *Beitr Pathol* 150:357–377
- Silverman SH, Migeon C, Rosemberg E, Wilkins L 1952 Precocious growth of sexual hair without other secondary sexual development: "premature pubarche," a constitutional variation of adolescence. *Pediatrics* 10:426–432
- Sigurjonsdottir TJ, Hayles AB 1968 Premature pubarche. *Clin Pediatr (Phila)* 7:29–33
- Pang S 1981 Precocious thelarche and premature adrenarche predictors. *Pediatr Ann* 10:340–345
- Pang S 1984 Premature adrenarche. *Pediatr Adolesc Endocrinol* 13:173–184
- Miller WL, Auchus RJ 2011 The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr Rev* 32:81–151
- Ghizzoni L, Milani S 2000 The natural history of premature adrenarche. *J Pediatr Endocrinol Metab* 13:1247–1251
- Ibáñez L, Virdis R, Potau N, Zampolli M, Ghizzoni L, Albisu MA, Carrascosa A, Bernasconi S, Vicens-Calvet E 1992 Natural history of premature pubarche: an auxological study. *J Clin Endocrinol Metab* 74:254–257
- Pere A, Perheentupa J, Peter M, Voutilainen R 1995 Follow up of growth and steroids in premature adrenarche. *Eur J Pediatr* 154:346–352
- Ibáñez L, Potau N, Marcos MV, de Zegher F 1999 Exaggerated adrenarche and hyperinsulinism in adolescent girls born small for gestational age. *J Clin Endocrinol Metab* 84:4739–4741
- Diaz A, Bhandari S, Sison C, Vogiatzi M 2008 Characteristics of children with premature pubarche in the New York metropolitan area. *Horm Res* 70:150–154
- Ibáñez L, Jiménez R, de Zegher F 2006 Early puberty-menarche after precocious pubarche: relation to prenatal growth. *Pediatrics* 117:117–121
- Denburg MR, Silfen ME, Manibo AM, Chin D, Levine LS, Ferin M, McMahon DJ, Go C, Oberfield SE 2002 Insulin sensitivity and the insulin-like growth factor system in prepubertal boys with premature adrenarche. *J Clin Endocrinol Metab* 87:5604–5609
- Oppenheimer E, Linder B, DiMartino-Nardi J 1995 Decreased insulin sensitivity in prepubertal girls with premature adrenarche and acanthosis nigricans. *J Clin Endocrinol Metab* 80:614–618
- Vuguin P, Linder B, Rosenfeld RG, Saenger P, DiMartino-Nardi J 1999 The roles of insulin sensitivity, insulin-like growth factor I (IGF-1), and IGF-binding protein-1 and -3 in the hyperandrogenism of African-American and Caribbean Hispanic girls with premature adrenarche. *J Clin Endocrinol Metab* 84:2037–2042
- Ibáñez L, Potau N, Zampolli M, Riqué S, Saenger P, Carrascosa A 1997 Hyperinsulinemia and decreased insulin-like growth factor-binding protein-1 are common features in prepubertal and pubertal girls with a history of premature pubarche. *J Clin Endocrinol Metab* 82:2283–2288
- Silfen ME, Manibo AM, Ferin M, McMahon DJ, Levine LS, Oberfield SE 2002 Elevated free IGF-1 levels in prepubertal Hispanic girls with premature adrenarche: relationship with hyperandrogenism and insulin sensitivity. *J Clin Endocrinol Metab* 87:398–403
- Ibáñez L, Aulesa C, Potau N, Ong K, Dunger DB, de Zegher F 2002 Plasminogen activator inhibitor-1 in girls with precocious pubarche: a premenarcheal marker for polycystic ovary syndrome? *Pediatr Res* 51:244–248
- Centers for Disease Control and Prevention National Center for Health

- Statistics 2008 National Health and Nutrition Examination Survey Data. Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. <http://www.cdc.gov/growthcharts/>
20. Silfen ME, Manibo AM, McMahon DJ, Levine LS, Murphy AR, Oberfield SE 2001 Comparison of simple measures of insulin sensitivity in young girls with premature adrenarche: the fasting glucose to insulin ratio may be a simple and useful measure. *J Clin Endocrinol Metab* 86:2863–2868
  21. Fernández JR, Redden DT, Pietrobelli A, Allison DB 2004 Waist circumference percentiles in nationally representative samples of African-American, European-American, and Mexican-American children and adolescents. *J Pediatr* 145:439–444
  22. Ohki Y, Orimo H, Kishi M, Ohkawa T 2004 Indexes of insulin resistance using the oral glucose tolerance test (O-GTT) in Japanese children and adolescents. *J Nippon Med Sch* 71:84–87
  23. McCarthy HD, Cole TJ, Fry T, Jebb SA, Prentice AM 2006 Body fat reference curves for children. *Int J Obes (Lond)* 30:598–602
  24. Lanman JT 1953 The fetal zone of the adrenal gland: its developmental course, comparative anatomy, and possible physiologic functions. *Medicine (Baltimore)* 32:389–430
  25. Mesiano S, Katz SL, Lee JY, Jaffe RB 1997 Insulin-like growth factors augment steroid production and expression of steroidogenic enzymes in human fetal adrenal cortical cells: implications for adrenal androgen regulation. *J Clin Endocrinol Metab* 82:1390–1396
  26. Spencer SJ, Mesiano S, Lee JY, Jaffe RB 1999 Proliferation and apoptosis in the human adrenal cortex during the fetal and perinatal periods: implications for growth and remodeling. *J Clin Endocrinol Metab* 84:1110–1115
  27. Hammer GD, Parker KL, Schimmer BP 2005 Minireview: transcriptional regulation of adrenocortical development. *Endocrinology* 146:1018–1024
  28. Stocco DM, Clark BJ 1996 Regulation of the acute production of steroids in steroidogenic cells. *Endocr Rev* 17:221–244
  29. Miller WL 2007 Steroidogenic acute regulatory protein (StAR), a novel mitochondrial cholesterol transporter. *Biochim Biophys Acta* 1771:663–676
  30. Nakamura Y, Gang HX, Suzuki T, Sasano H, Rainey WE 2009 Adrenal changes associated with adrenarche. *Rev Endocr Metab Disord* 10:19–26
  31. Auchus RJ, Rainey WE 2004 Adrenarche—physiology, biochemistry and human disease. *Clin Endocrinol (Oxf)* 60:288–296
  32. Miller WL, Auchus RJ, Geller DH 1997 The regulation of 17,20 lyase activity. *Steroids* 62:133–142
  33. Miller WL, Huang N, Pandey AV, Flück CE, Agrawal V 2005 P450 oxidoreductase deficiency: a new disorder of steroidogenesis. *Ann NY Acad Sci* 1061:100–108
  34. Lin D, Black SM, Nagahama Y, Miller WL 1993 Steroid 17 $\alpha$ -hydroxylase and 17,20-lyase activities of P450c17: contributions of serine106 and P450 reductase. *Endocrinology* 132:2498–2506
  35. Auchus RJ, Geller DH, Lee TC, Miller WL 1998 The regulation of human P450c17 activity: relationship to premature adrenarche, insulin resistance and the polycystic ovary syndrome. *Trends Endocrinol Metab* 9:47–50
  36. Mapes S, Corbin CJ, Tarantal A, Conley A 1999 The primate adrenal zona reticularis is defined by expression of cytochrome b5, 17 $\alpha$ -hydroxylase/17,20-lyase cytochrome P450 (P450c17) and NADPH-cytochrome P450 reductase (reductase) but not 3 $\beta$ -hydroxysteroid dehydrogenase/ $\Delta$ 5–4 isomerase (3 $\beta$ -HSD). *J Clin Endocrinol Metab* 84:3382–3385
  37. Suzuki T, Sasano H, Takeyama J, Kaneko C, Freije WA, Carr BR, Rainey WE 2000 Developmental changes in steroidogenic enzymes in human postnatal adrenal cortex: immunohistochemical studies. *Clin Endocrinol (Oxf)* 53:739–747
  38. Zhang LH, Rodriguez H, Ohno S, Miller WL 1995 Serine phosphorylation of human P450c17 increases 17,20-lyase activity: implications for adrenarche and the polycystic ovary syndrome. *Proc Natl Acad Sci USA* 92:10619–10623
  39. Pandey AV, Mellon SH, Miller WL 2003 Protein phosphatase 2A and phosphoprotein SET regulate androgen production by P450c17. *J Biol Chem* 278:2837–2844
  40. Chin JE, Dickens M, Tavare JM, Roth RA 1993 Overexpression of protein kinase C isoenzymes  $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\epsilon$  in cells overexpressing the insulin receptor. Effects on receptor phosphorylation and signaling. *J Biol Chem* 268:6338–6347
  41. Takayama S, White MF, Kahn CR 1988 Phorbol ester-induced serine phosphorylation of the insulin receptor decreases its tyrosine kinase activity. *J Biol Chem* 263:3440–3447
  42. Lorence MC, Murry BA, Trant JM, Mason JI 1990 Human 3 $\beta$ -hydroxysteroid dehydrogenase/ $\Delta$ 5–4 isomerase from placenta: expression in non-steroidogenic cells of a protein that catalyzes the dehydrogenation/isomerization of C21 and C19 steroids. *Endocrinology* 126:2493–2498
  43. Topor LS, Asai M, Dunn J, Majzoub JA 2011 Cortisol stimulates secretion of dehydroepiandrosterone in human adrenocortical cells through inhibition of 3 $\beta$ HSD2. *J Clin Endocrinol Metab* 96:E31–E39
  44. Gell JS, Carr BR, Sasano H, Atkins B, Margraf L, Mason JI, Rainey WE 1998 Adrenarche results from development of a 3 $\beta$ -hydroxysteroid dehydrogenase-deficient adrenal reticularis. *J Clin Endocrinol Metab* 83:3695–3701
  45. Noordam C, Dhir V, McNelis JC, Schlereth F, Hanley NA, Krone N, Smeitink JA, Smeets R, Sweep FC, Claahsen-van der Grinten HL, Arlt W 2009 Inactivating PAPS2 mutations in a patient with premature pubarche. *N Engl J Med* 360:2310–2318
  46. Duncan GE, Li SM, Zhou XH 2004 Prevalence and trends of a metabolic syndrome phenotype among US adolescents, 1999–2000. *Diabetes Care* 27:2438–2443
  47. Weiss R, Dziura J, Burgert TS, Tamborlane WV, Taksali SE, Yeckel CW, Allen K, Lopes M, Savoye M, Morrison J, Sherwin RS, Caprio S 2004 Obesity and the metabolic syndrome in children and adolescents. *N Engl J Med* 350:2362–2374
  48. Ibáñez L, Potau N, Chacon P, Pascual C, Carrascosa A 1998 Hyperinsulinaemia, dyslipaemia and cardiovascular risk in girls with a history of premature pubarche. *Diabetologia* 41:1057–1063
  49. Ibáñez L, Castell C, Tresserras R, Potau N 1999 Increased prevalence of unknown type 2 diabetes mellitus and impaired glucose tolerance in first-degree relatives of girls with a history of precocious pubarche. *Clin Endocrinol (Oxf)* 51:395–401
  50. Cara JF, Rosenfield RL 1988 Insulin-like growth factor I and insulin potentiate luteinizing hormone-induced androgen synthesis by rat ovarian thecal-interstitial cells. *Endocrinology* 123:733–739
  51. Duleba AJ, Spaczynski RZ, Olive DL 1998 Insulin and insulin-like growth factor I stimulate the proliferation of human ovarian theca-interstitial cells. *Fertil Steril* 69:335–340
  52. Plymate SR, Hoop RC, Jones RE, Matej LA 1990 Regulation of sex hormone-binding globulin production by growth factors. *Metabolism* 39:967–970
  53. Ibáñez L, Dimartino-Nardi J, Potau N, Saenger P 2000 Premature adrenarche—normal variant or forerunner of adult disease? *Endocr Rev* 21:671–696
  54. Rosenfield RL 1996 Editorial: evidence that idiopathic functional adrenal hyperandrogenism is caused by dysregulation of adrenal steroidogenesis and that hyperinsulinemia may be involved. *J Clin Endocrinol Metab* 81:878–880
  55. Dimartino-Nardi J 1999 Premature adrenarche: findings in prepubertal African-American and Caribbean-Hispanic girls. *Acta Paediatr Suppl* 88:67–72
  56. DiMartino-Nardi J 2000 Pre- and postpubertal findings in premature adrenarche. *J Pediatr Endocrinol Metab* 13(Suppl 5):1265–1269
  57. Vuguin P, Saenger P, Dimartino-Nardi J 2001 Fasting glucose insulin ratio: a useful measure of insulin resistance in girls with premature adrenarche. *J Clin Endocrinol Metab* 86:4618–4621
  58. McCartney CR, Prendergast KA, Chhabra S, Eagleson CA, Yoo R, Chang RJ, Foster CM, Marshall JC 2006 The association of obesity and hyperandrogenemia during the pubertal transition in girls: obesity as a potential factor in the genesis of postpubertal hyperandrogenism. *J Clin Endocrinol Metab* 91:1714–1722
  59. Ibáñez L, Potau N, Virdis R, Zampolli M, Terzi C, Gussinyé M, Carrascosa A, Vicens-Calvet E 1993 Postpubertal outcome in girls diagnosed of premature pubarche during childhood: increased frequency of functional ovarian hyperandrogenism. *J Clin Endocrinol Metab* 76:1599–1603
  60. Ibáñez L, Díaz R, López-Bermejo A, Marcos MV 2009 Clinical spectrum of premature pubarche: links to metabolic syndrome and ovarian hyperandrogenism. *Rev Endocr Metab Disord* 10:63–76
  61. Ibáñez L, Valls C, Potau N, Marcos MV, de Zegher F 2001 Polycystic ovary syndrome after precocious pubarche: ontogeny of the low-birth-weight effect. *Clin Endocrinol (Oxf)* 55:667–672
  62. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop

- Group 2004 Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril* 81:19–25
63. Zawadzki JK, Dunaif A 1992 Diagnostic criteria for polycystic ovary syndrome: towards a rational approach. In: Dunaif A, Givens JR, Haseltine FP, Merriam GR, eds. *Polycystic ovary syndrome*. Oxford, UK: Blackwell; 59–69
  64. Carmina E, Oberfield SE, Lobo RA 2010 The diagnosis of polycystic ovary syndrome in adolescents. *Am J Obstet Gynecol* 203:201.e1–5
  65. Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH 2003 Prevalence of a metabolic syndrome phenotype in adolescents: findings from the third National Health and Nutrition Examination Survey, 1988–1994. *Arch Pediatr Adolesc Med* 157:821–827
  66. de Ferranti SD, Gauvreau K, Ludwig DS, Neufeld EJ, Newburger JW, Rifai N 2004 Prevalence of the metabolic syndrome in American adolescents: findings from the Third National Health and Nutrition Examination Survey. *Circulation* 110:2494–2497
  67. Güven A, Cinaz P, Bideci A 2005 Is premature adrenarche a risk factor for atherogenesis? *Pediatr Int* 47:20–25
  68. Andiran N, Yordam N 2008 Lipoprotein(a) levels in girls with premature adrenarche. *J Paediatr Child Health* 44:138–142
  69. de Zegher F, Lopez-Bermejo A, Ibáñez L 2009 Adipose tissue expandability and the early origins of PCOS. *Trends Endocrinol Metab* 20:418–423
  70. Leibel N, Shen W, Mao X, Punyanitya M, Gallagher D, Horlick M, Shungu DC, Oberfield SE 2009 Body composition in premature adrenarche by structural MRI, 1H MRS and DXA. *J Pediatr Endocrinol Metab* 22:301–307
  71. Sopher AB, Thornton JC, Silfen ME, Manibo A, Oberfield SE, Wang J, Pierson Jr RN, Levine LS, Horlick M 2001 Prepubertal girls with premature adrenarche have greater bone mineral content and density than controls. *J Clin Endocrinol Metab* 86:5269–5272
  72. Utriainen P, Jääskeläinen J, Saarinen A, Vanninen E, Mäkitie O, Voutilainen R 2009 Body composition and bone mineral density in children with premature adrenarche and the association of LRP5 gene polymorphisms with bone mineral density. *J Clin Endocrinol Metab* 94:4144–4151
  73. Ibáñez L, Ong K, de Zegher F, Marcos MV, del Rio L, Dunger DB 2003 Fat distribution in non-obese girls with and without precocious pubarche: central adiposity related to insulinaemia and androgenaemia from prepuberty to postmenarche. *Clin Endocrinol (Oxf)* 58:372–379
  74. Ibáñez L, De Zegher F 2003 Flutamide-metformin therapy to reduce fat mass in hyperinsulinemic ovarian hyperandrogenism: effects in adolescents and in women on third-generation oral contraceptives. *J Clin Endocrinol Metab* 88:4720–4724
  75. Ibáñez L, Ferrer A, Ong K, Amin R, Dunger D, de Zegher F 2004 Insulin sensitization early post-menarche prevents progression from precocious pubarche to polycystic ovary syndrome. *J Pediatr* 144:23–29
  76. Ibáñez L, Ong K, Ferrer A, Amin R, Dunger D, de Zegher F 2003 Low-dose flutamide-metformin therapy reverses insulin resistance and reduces fat mass in nonobese adolescents with ovarian hyperandrogenism. *J Clin Endocrinol Metab* 88:2600–2606
  77. Ibáñez L, Potau N, Ferrer A, Rodriguez-Hierro F, Marcos MV, De Zegher F 2002 Anovulation in eumenorrheic, nonobese adolescent girls born small for gestational age: insulin sensitization induces ovulation, increases lean body mass, and reduces abdominal fat excess, dyslipidemia, and subclinical hyperandrogenism. *J Clin Endocrinol Metab* 87:5702–5705
  78. Vottero A, Capelletti M, Giuliodori S, Viani I, Ziveri M, Neri TM, Bernasconi S, Ghizzoni L 2006 Decreased androgen receptor gene methylation in premature pubarche: a novel pathogenetic mechanism? *J Clin Endocrinol Metab* 91:968–972
  79. Kaplan SL, Grumbach MM 1998 Pathogenesis of sexual precocity. In: Grumbach MM, Sizonenko PC, Aubert ML, eds. *Williams textbook of endocrinology*. Philadelphia: WB Saunders; 1509–1625
  80. Chalumeau M, Chemaitilly W, Trivin C, Adan L, Bréart G, Brauner R 2002 Central precocious puberty in girls: an evidence-based diagnosis tree to predict central nervous system abnormalities. *Pediatrics* 109:61–67
  81. Herman-Giddens ME, Slora EJ, Wasserman RC, Bourdony CJ, Bhappkar MV, Koch GG, Hasemeier CM 1997 Secondary sexual characteristics and menses in young girls seen in office practice: a study from the Pediatrics Research in Office Settings network. *Pediatrics* 99:505–512
  82. Wu T, Mendola P, Buck GM 2002 Ethnic differences in the presence of secondary sex characteristics and menarche among US girls: the third National Health and Nutrition Examination Survey, 1988–1994. *Pediatrics* 110:752–757
  83. Thamdrup E 1961 Precocious sexual development: a clinical study of 100 patients. Springfield, IL: Charles C. Thomas Publisher Ltd.; 50
  84. Paul D, Conte FA, Grumbach MM, Kaplan SL 1995 Long-term effect of gonadotropin-releasing hormone agonist therapy on final and near-final height in 26 children with true precocious puberty treated at a median age of less than 5 years. *J Clin Endocrinol Metab* 80:546–551
  85. Pescovitz OH, Cassorla F, Comite F, Loriaux DL, Cutler Jr GB 1985 LHRH analog treatment of central precocious puberty complicating congenital adrenal hyperplasia. *Ann NY Acad Sci* 458:174–181
  86. Pescovitz OH, Hench K, Green O, Comite F, Loriaux DL, Cutler Jr GB 1985 Central precocious puberty complicating a virilizing adrenal tumor: treatment with a long-acting LHRH analog. *J Pediatr* 106:612–614
  87. Carel JC, Léger J 2008 Precocious puberty. *N Engl J Med* 358:2366–2377
  88. Kohn B, Levine LS, Pollack MS, Pang S, Lorenzen F, Levy D, Lerner AJ, Rondanini GF, Dupont B, New MI 1982 Late-onset steroid 21-hydroxylase deficiency: a variant of classical congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 55:817–827
  89. Armengaud JB, Charkaluk ML, Trivin C, Tardy V, Bréart G, Brauner R, Chalumeau M 2009 Precocious pubarche: distinguishing late-onset congenital adrenal hyperplasia from premature adrenarche. *J Clin Endocrinol Metab* 94:2835–2840
  90. Voelker R 2010 Estrogen spray poses risks to children, pets through contact with treated skin. *JAMA* 304:953
  91. Henley DV, Lipson N, Korach KS, Bloch CA 2007 Prepubertal gynecostasia linked to lavender and tea tree oils. *N Engl J Med* 356:479–485
  92. Vandenberg LN, Maffini MV, Sonnenschein C, Rubin BS, Soto AM 2009 Bisphenol-A and the great divide: a review of controversies in the field of endocrine disruption. *Endocr Rev* 30:75–95
  93. Kunz GJ, Klein KO, Clemons RD, Gottschalk ME, Jones KL 2004 Virilization of young children after topical androgen use by their parents. *Pediatrics* 114:282–284
  94. Diamanti-Kandarakis E, Economou F, Palimeri S, Christakou C 2010 Metformin in polycystic ovary syndrome. *Ann NY Acad Sci* 1205:192–198
  95. Palomba S, Falbo A, Zullo F, Orio Jr F 2009 Evidence-based and potential benefits of metformin in the polycystic ovary syndrome: a comprehensive review. *Endocr Rev* 30:1–50
  96. Ibáñez L, López-Bermejo A, Diaz M, Marcos MV, de Zegher F 2008 Metformin treatment for four years to reduce total and visceral fat in low birth weight girls with precocious pubarche. *J Clin Endocrinol Metab* 93:1841–1845
  97. Ibáñez L, Lopez-Bermejo A, Diaz M, Marcos MV, de Zegher F 2011 Early metformin therapy to delay menarche and augment height in girls with precocious pubarche. *Fertil Steril* 95:727–730
  98. Ibáñez L, Lopez-Bermejo A, Diaz M, Marcos MV, de Zegher F 2010 Pubertal metformin therapy to reduce total, visceral, and hepatic adiposity. *J Pediatr* 156:98–102.e1
  99. Euling SY, Herman-Giddens ME, Lee PA, Selevan SG, Juul A, Sorensen TI, Dunkel L, Himes JH, Teilmann G, Swan SH 2008 Examination of US puberty-timing data from 1940 to 1994 for secular trends: panel findings. *Pediatrics* 121:S172–191
  100. Petersen KF, Dufour S, Savage DB, Bilz S, Solomon G, Yonemitsu S, Cline GW, Befroy D, Zeman L, Kahn BB, Papademetris X, Rothman DL, Shulman GI 2007 The role of skeletal muscle insulin resistance in the pathogenesis of the metabolic syndrome. *Proc Natl Acad Sci USA* 104:12587–12594
  101. Sopher AB, Lee E, Root J, Castle RH, Frisse EA, Gallagher D, Hirsch J, Mao X, Shungu DK, Hassoun A, McMahon DJ, Oberfield SE, Intramyocellular lipid (IMCL) deposition is an early marker of insulin resistance (IR) in adolescent girls with polycystic ovary syndrome (PCOS) and controls. Program of the Annual Meeting of the Pediatric Academic Societies, Vancouver, British Columbia, 2010. (Abstract 2855.402)
  102. Ibáñez L, de Zegher F 2004 Ethinylestradiol-drospirenone, flutamide-metformin, or both for adolescents and women with hyperinsulinemic hyperandrogenism: opposite effects on adipocytokines and body adiposity. *J Clin Endocrinol Metab* 89:1592–1597
  103. National Heart, Lung, and Blood Institute 1980 The Lipid Research Clinics Population Studies Data Book. Vol. I. The prevalence study. Bethesda, MD: U.S. Department of Heart and Human Services, Public Health Service, National Institutes of Health, NIH pub. no. 80-1527